



PATIENT NAME:

DOB:

PHARMACY:

POSTOPERATIVE INSTRUCTIONS

PROCEDURE:

NEXT APPOINTMENT:

LOCATION:

PAIN MEDICATION

You will be given a prescription for pain medication after surgery. This should be filled and ready for use when you return home from surgery. You **SHOULD NOT** drive, operate heavy machinery, or participate in activities that require concentration while taking narcotic pain medications such as Percocet, Oxycodone, Norco, Vicodin, or Tylenol with Codeine. You **SHOULD** take an over-the-counter stool softener while taking narcotic pain medications to prevent constipation.

DRESSING

- DO NOT REMOVE.** Keep dressing clean and dry **AT ALL TIMES**
- Dressing may be removed **72 HOURS AFTER SURGERY** and begin dry/sterile dressing changes. Keep incisions covered, clean, and dry. **DO NOT** get incision(s) wet or soak them in any water.

WEIGHTBEARING STATUS

- WEIGHTBEARING AS TOLERATED** with crutches
- NONWEIGHTBEARING** with crutches

COLD THERAPY MACHINE

If provided, begin using your cold therapy machine after surgery. We recommend use at least 3 times per day, following the manufacturers instructions for setup and utilization of water and ice, for **NO LONGER THAN 45 MINUTES** per treatment. **ALWAYS** make sure to apply a barrier between the cooling pad and your skin. If you have not been provided with a cold therapy machine, apply ice to the surgical area in a waterproof bag for **NO LONGER THAN 30 MINUTES** per treatment.

DVT (BLOOD CLOT) PREVENTION

- >> Beginning the day of surgery, take one **ASPIRIN 81MG** tablet, twice per day for 4 weeks unless not allowed for other medical reasons. **Clear this with your family doctor before starting the aspirin.**
- >> Perform ankle range of motion exercises (ankle pumps) on uninjured leg to exercise calf muscles. Do ten reps every hour minimum while awake.
- >> **You should not do any flying for 2 weeks after surgery**

PAIN MEDICATION REFILL PROTOCOL

You **MUST** give your physician **48 HOURS** notice for any medication refill. Many medications require your surgeon to send the medication electronically and cannot be called into the pharmacy. Prescriptions **WILL NOT** be filled over the weekend. A prescription for the same medication you have been taking after surgery **IS NOT GUARANTEED.**

WHEN TO CALL YOUR SURGEON

Complications after surgery are fortunately very rare. Please contact our office if you have any of the following symptoms:

- >> Persistent fever >101°F, chills
- >> Increasing pain at the surgical site

- >> Increased swelling/pain in the calf
- >> New onset numbness or tingling
- >> Hives or new rashes
- >> Shortness of breath or chest pain
- >> Persistent nausea/vomiting
- >> Drainage from your incision lasting >5 days